

Accepted Manuscript


An unusual silver-ethynide polymeric chain containing centrosymmetric Ag₁₄ cluster segments stabilized by mixed carboxylate ligands

Yun-Peng Xie, Shael A. Al-Thabaiti, Mohamed Mokhtar, Thomas C.W. Mak

PII: S1387-7003(13)00088-9
DOI: doi: [10.1016/j.inoche.2013.02.019](https://doi.org/10.1016/j.inoche.2013.02.019)
Reference: INOCHE 4992

To appear in: *Inorganic Chemistry Communications*

Received date: 26 January 2013
Accepted date: 27 February 2013



Please cite this article as: Yun-Peng Xie, Shael A. Al-Thabaiti, Mohamed Mokhtar, Thomas C.W. Mak, An unusual silver-ethynide polymeric chain containing centrosymmetric Ag₁₄ cluster segments stabilized by mixed carboxylate ligands, *Inorganic Chemistry Communications* (2013), doi: [10.1016/j.inoche.2013.02.019](https://doi.org/10.1016/j.inoche.2013.02.019)

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

An unusual silver-ethynide polymeric chain containing centrosymmetric Ag₁₄ cluster segments stabilized by mixed carboxylate ligands

Yun-Peng Xie,^{a,b} Shaeel A. Al-Thabaiti,^c Mohamed Mokhtar,^c Thomas C. W. Mak^{a,c*}

^a Department of Chemistry and Center of Novel Functional Molecules, The Chinese University of Hong Kong, Shatin, New Territories, Hong Kong SAR, People's Republic of China

^b College of Materials Science and Engineering, Huazhong University of Science and Technology, Wuhan, Hubei Province, People's Republic of China

^c Department of Chemistry, King Abdulaziz Univeristy, Jeddah 21589, Kingdom of Saudi Arabia

*Author for correspondence

Tel: (852) 3943 6279

Fax: (852) 2603 5057

E-mail: tcwmak@cuhk.edu.hk

Abstract

From the reaction of AgC≡C^tBu with 9-hydroxy-9-fluorenicarboxylic acid (HL) and AgCF₃CO₂, we have isolated a new type of organosilver(I) coordination polymeric chain, namely $\{[Ag_{14}(C\equiv C^tBu)_8(L)_2(CF_3CO_2)_4]\cdot 2CH_3OH\}_n$, which contains centrosymmetric $[Ag_{14}(C\equiv C^tBu)_8(CF_3CO_2)_2]^{4+}$ cluster segments bridged by pairs of L and trifluoroacetate ligands.

Keywords: metal-ligand supramolecular synthon; organosilver complex; polymeric coordination chain; silver(I) ethynide cluster

The chemistry of metal ethynyl complexes is of great interest because of their structural diversity and technological application as precursors of nonlinear optical materials, luminescent probes, and rigid-rod molecular wires [1-4]. Among a wide range of candidate metals, the Group 11 elements (Cu, Ag and Au) are frequently employed in the syntheses of novel ethynyl complexes due to their enhanced metallophilic interaction that overcomes repulsion between cationic metal centers [5-11]. The silver(I) ethynyl derivatives are less explored because of their sensitivity to light and tendency to yield highly insoluble polymers/oligomers. Recently, we and other groups have prepared and structurally characterized a series of silver(I)-ethynide coordination polymeric networks [12-17] and high-nuclearity clusters [18-31] based on the multinuclear metal-ligand supramolecular synthon $R-C\equiv C\supset Ag_n$ ($n = 3, 4, 5$; R = alkyl, aryl, heteroaryl).

In 1999 the reaction of $[AgC\equiv C^tBu]_n$ with $AgBF_4$ in molar ratio 2:1 was reported to yield a polymeric undulated ribbon structure $[Ag_3(C\equiv C^tBu)_2^+]_n$ [32]. Envisioning that such soluble species might be good precursors for the preparation of new silver(I)-ethynide complexes, we have developed a synthetic strategy by introducing phosphonate and phosphonate-functionalized oxovanadate or oxovanadate as capping ligands for trapping oligomeric silver(I)-ethynide clusters. Recently we isolated a series of mixed-metal giant clusters with the silver(I)-*tert*-butylethynide moiety and various phosphonate-functionalized oxovanadate building blocks as their surface components, as well as different numbers and kinds of anionic species as their encapsulated templates: for example $\{(NO_3)_2@Ag_{16}(C\equiv CPh)_4[(^tBuPO_3)_4V_4O_8]_2(DMF)_6(NO_3)_2\}$ [31] and $\{Cl_2@Ag_{21}(C\equiv C^tBu)_9[(^tBuPO_3)_3V_3O_6(OH)]_2[(^tBuPO_3)VO_2(OH)](MeOH)_2(H_2O)_2\}$ [30]. We also employed neat *tert*-butylphosphonic acid as a precursor and successfully generated new silver(I) ethynide networks and clusters, for example $3AgC\equiv CPh\cdot Ag_2^tBuPO_3\cdot Ag^tBuPO_3H\cdot 2AgNO_3$ and $[Ag_8(Cl@Ag_{14})](C\equiv C^tBu)_{14}(^tBuPO_3)_2F_2(H_2O)_2]BF_4\cdot 3.5H_2O$ [16].

In the present work, we report our successful combination of 9-hydroxy-9-fluorene-carboxylic acid (HL) with the supramolecular synthon $^tBuC\equiv C\supset Ag_n$ (n

= 3, 4) to generate a novel organosilver coordination polymer, namely $\{[\text{Ag}_{14}(\text{C}\equiv\text{C}^t\text{Bu})_8(\text{L})_2(\text{CF}_3\text{CO}_2)_4]\cdot 2\text{CH}_3\text{OH}\}_n$ (**1**) [33], which has been characterized by elemental analysis, infrared spectroscopy and single-crystal X-ray analysis [34]. HL is known to function as a plant growth regulator, which restrains the generation of abscisic acid and regulate plant root calcium transport [35]. Furthermore, in the anionic L ligand, the hydroxy and carboxylate groups provide different potential coordination sites for metal coordination.

The reaction of HL with $\text{AgC}\equiv\text{C}^t\text{Bu}$ and AgCF_3CO_2 in CH_3OH yielded colorless prismatic crystals of **1**. An IR vibration band at 2005 cm^{-1} confirmed the presence of the $\text{C}\equiv\text{C}$ functional group. Of the four independent *tert*-butylethyne ligands in the asymmetric unit, the ethynide group composed of C37 and C38 is bound to a silver triangle in the $\mu_3\text{-}\eta^1, \eta^1, \eta^1$ coordination mode. The remaining three ethynide groups (C15 \equiv C16, C23 \equiv C24 and C31 \equiv C32) are each capped by a butterfly-shaped Ag_4 basket in the $\mu_4\text{-}\eta^1, \eta^1, \eta^1, \eta^2$ or $\mu_4\text{-}\eta^1, \eta^1, \eta^2, \eta^2$ coordination mode, as shown in Fig. 1.

Figure 1 here

Two Ag_3 triangles and six Ag_4 baskets coalesce by sharing vertices to form a centrosymmetric $\text{Ag}_{14}(\text{}^t\text{BuC}\equiv\text{C})_8$ cluster, which is further stabilized by a pair of CF_3CO_2 ligands with $\text{Ag}\text{-O}$ bond distances of 2.317(5) and 2.369(6) Å (Fig. 2). The resulting $\text{Ag}_{14}(\text{}^t\text{BuC}\equiv\text{C})_8(\text{CF}_3\text{CO}_2)_2$ cluster segments are bridged by a series of L and additional CF_3CO_2 ligands to engender an infinite coordination chain in the direction of the *a* axis (Fig. 3). The L ligand has its carboxylate terminal bridging two adjacent silver atoms, and the 9-hydroxy group functions as a monodentate ligand to coordinate one silver atom, with $\text{Ag}\text{-O}$ distances varying from 2.343(6) to 2.589(5) Å. However, the bridging CF_3CO_2 ligand coordinates to four silver atoms with $\text{Ag}\text{-O}$ bond lengths ranging from 2.335(6) to 2.715(6) Å.

Figure 2 here

Figure 3 here

The argentophilic Ag...Ag bond distances lie in the range 2.899(2)–3.325(2) Å. Additionally, the crystal structure contains two methanol solvate molecules in the unit cell. Fig. 4 shows the crystal structure with only van der Waals interaction between the coordination polymeric chains.

Figure 4 here

Notably, the bar-like Ag₁₄ cluster in **1** differs markedly from previously reported rhombic-dodecahedral silver(I)-ethynide cages, e.g. [Cl@Ag₁₄(C≡C^tBu)₁₂]OH [19] and [Cl@Ag₁₄(C≡C_{cyclohexyl})₁₂]Cl [28] with encapsulated chloride ions, which have been employed as precursors to synthesize the corresponding iso-structural “cluster-within-cluster” compounds [Cl₆Ag₈@Ag₃₀(^tBuC≡C)₂₀(ClO₄)₁₂]·Et₂O and [Cl₆Ag₈@Ag₃₀(chxC≡C)₂₀(ClO₄)₁₀](ClO₄)₂·1.5Et₂O that bear the same pseudo-*O_h* cationic Cl₆Ag₈ inner core [28].

The anionic L ligand, with its bulky hydrophobic, outward-extending fluorenyl group, connects adjacent high-nuclearity silver(I)-ethynide cluster segments to generate the coordination polymer chain in **1**. In contrast, silver trifluoroacetate plays a crucial dual role in the present synthesis: (i) as a bridging ligand that stabilizes each Ag₁₄ cluster segment, and (ii) cooperates with the L ligand to link up adjacent Ag₁₄ cluster segments. Notably, thus far our attempts to prepare related silver(I)-ethynide complexes using other soluble silver salts such as nitrate, tetrafluoroborate and triflate were unsuccessful, which always yielded an intractable pale solid residue.

Conclusion

Combining the metal-ligand supramolecular synthon ^tBuC≡C⊃Ag_{*n*} (*n* = 3, 4) (generated *in situ* from a solution of ^tBuC≡CAg and AgCF₃CO₂ in methanol) and 9-hydroxy-9-fluorenicarboxylic acid, we have isolated and structurally characterized a new kind of silver(I)-ethynide coordination polymeric complex, in which the 9-hydroxy-9-fluorenicarboxylate and trifluoroacetate ligands bridge two novel condensed

Ag₁₄ clusters to form a coordination chain architecture. The present study offers broad prospects to future synthetic exploration of silver-ethynide complexes bearing different varieties of carboxylates as structural components.

Acknowledgments

This work was supported by the Wei Lun Foundation, Grant No. 27-3-1432/HiCi of King Abdulaziz University, and a Postdoctoral Research Fellowship award to Y.-P. Xie by The Chinese University of Hong Kong.

Appendix A. Supplementary material

CCDC 907576 contains the supplementary crystallographic data for this paper, which can be obtained free of charge from The Cambridge Crystallographic Data Centre via www.ccdc.cam.ac.uk/data_request/cif.

References

- [1] M.I. Bruce, Organometallic chemistry of vinylidene and related unsaturated carbenes *Chem. Rev.* 91 (1991) 197–257.
- [2] F. Paul, C. Lapinte, Organometallic molecular wires and other nanoscale-sized devices: An approach using the organoiron (dppe)Cp*Fe building block, *Coord. Chem. Rev.* 178–180 (1998) 431–509.
- [3] V.W.–W. Yam, Molecular Design of Transition Metal Alkynyl Complexes as Building Blocks for Luminescent Metal-Based Materials: Structural and Photophysical Aspects, *Acc. Chem. Res.* 35 (2002) 555–563.
- [4] V.W.–W. Yam, E.C.–C. Cheng, Highlights on the recent advances in gold chemistry—a photophysical perspective, *Chem. Soc. Rev.* 37 (2008) 1806–1813.
- [5] M. Jansen, Homoatomic d¹⁰–d¹⁰ Interactions: Their Effects on Structure and Chemical and Physical Properties, *Angew. Chem. Int. Ed.* 26 (1987) 1098–1110.

- [6] P. Pyykkö, Strong Closed-Shell Interactions in Inorganic Chemistry, *Chem. Rev.* 97 (1997) 597-636.
- [7] P. Pyykkö, Theoretical Chemistry of Gold, *Angew. Chem. Int. Ed.* 43 (2004) 4412-4456.
- [8] V.W.-W. Yam, K.M.-C. Wong, Luminescent Molecular Rods-Transition-Metal Alkynyl Complexes, *Top. Curr. Chem.* 257 (2005) 1-32.
- [9] R.J. Puddephatt, Macrocycles, catenanes, oligomers and polymers in gold chemistry, *Chem. Soc. Rev.* 37 (2008) 2012-2027.
- [10] H.L. Hermann, G. Boche, P. Schwerdtfeger, Metallophilic Interactions in Closed-Shell Copper(I) Compounds—A Theoretical Study, *Chem. Eur. J.* 7 (2001) 5333-5342.
- [11] C.J. Adams, M.I. Bruce, E. Horn, B.W. Skelton, E.R.T. Tiekink, A.H. White, Cluster chemistry. Part 91. Clusters derived from 1,4-Bis (diphenylphosphino)buta-1,3-diyne and their pyrolysis products: a route to complexes containing the tetracarbon ligand, *J. Chem. Soc. Dalton Trans.* (1993) 3299-3312.
- [12] L. Zhao, W.-Y. Wong, T.C.W. Mak, Novel μ_5 -Coordination Modes of Aryl and Alkyl Ethynides and Classification of Metal-Ligand Interactions in Silver(I) Complexes, *Chem. Eur. J.* 12 (2006) 4865-4872.
- [13] S.-Q. Zang, J. Han, T.C.W. Mak, Silver(I)-Organic Networks Assembled with the Flexible Prop-2-ynoxybenzene Ligand: In Situ Recrystallization and Unusual Silver-Aromatic Interaction, *Organometallics* 28 (2009) 2677-2683.
- [14] B. Li, S.-Q. Zang, R. Liang, Y.-J. Wu, T.C.W. Mak, Silver(I)-Organic Networks Assembled with Propargyl-Functionalized Di- and Trihydroxybenzenes, *Organometallics* 30 (2011) 1710-1718.
- [15] Y. Zhao, P. Zhang, B. Li, X. G. Meng, T.L. Zhang, Two- and Three-Dimensional Silver(I)-Organic Networks Generated from Mono- and Dicarboxylphenylethyne, *Inorg. Chem.* 50 (2011), 9097-9105.
- [16] Y.-P. Xie, T.C.W. Mak, Silver(I) Ethynide Coordination Networks and Clusters Assembled with *tert*-Butylphosphonic Acid, *Inorg. Chem.* 51 (2012) 8640-8642.

- [17] P.-S. Cheng, S. Marivel, S.-Q. Zang, G.-G. Gao, T.C.W. Mak, Argentophilic Infinite Chain, Column, and Layer Structures Assembled with the Multinuclear Silver(I)–Phenylethyne Supramolecular Synthons, *Cryst. Growth Des.* 12 (2012) 4519–4529.
- [18] D. Rais, J. Yau, D.M.P. Mingos, R. Vilar, A.J.P. White, D.J. Williams, Anion-Templated Syntheses of Rhombohedral Silver–Alkynyl Cage Compounds, *Angew. Chem. Int. Ed.* 40 (2001) 3464–3467.
- [19] D. Rais, D.M.P. Mingos, R. Vilar, A.J.P. White, D.J. Williams, Directing role of anions in the syntheses of the silver-alkynyl cages $[\text{Ag}_{14}(\text{C}\equiv\text{C}^t\text{Bu})_{12}\text{X}][\text{BF}_4]$ (X=F, Cl, Br) and silver-alkynyl polymers $[\text{Ag}_3(\text{C}\equiv\text{C}^t\text{Bu})_2(\text{X})]_n$ (X=Tos, NO_3), *J. Organomet. Chem.* 652 (2002) 87–93.
- [20] S.-D. Bian, H.-B. Wu, Q.-M. Wang, A Facile Template Approach to High-Nuclearity Silver(I) Alkynyl Clusters, *Angew. Chem. Int. Ed.* 48 (2009) 5363–5365.
- [21] S.-D. Bian, J.-H. Jia, Q.-M. Wang, High-Nuclearity Silver Clusters Templated by Carbonates Generated from Atmospheric Carbon Dioxide Fixation, *J. Am. Chem. Soc.* 131, (2009) 3422–3423.
- [22] J. Qiao, K. Shi, Q.-M. Wang, A Giant Silver Alkynyl Cage with Sixty Silver(I) Ions Clustered around Polyoxometalate Templates, *Angew. Chem. Int. Ed.* 49 (2010) 1765–1767.
- [23] P. Putaj, F. Lefebvre, Polyoxometalates containing late transition and noble metal atoms, *Coord. Chem. Rev.* 255 (2011) 1642–1685.
- [24] C.-Y. Gao, L. Zhao, M.-X. Wang, Designed Synthesis of Metal Cluster-Centered Pseudo-Rotaxane Supramolecular Architectures, *J. Am. Chem. Soc.* 133 (2011) 8448–8451.
- [25] F. Gruber, M. Jansen, $\{[\text{Ag}_{42}(\text{CO}_3)(\text{C}\equiv\text{C}^t\text{Bu})_{27}(\text{CH}_3\text{CN})_2][\text{CoW}_{12}\text{O}_{40}]_2\}[\text{BF}_4]$: An Intercluster Sandwich Compound, *Angew. Chem. Int. Ed.* 49 (2010) 4924–4926.

- [26] G.-G. Gao, P.-S. Cheng, T.C.W. Mak, Acid-Induced Surface Functionalization of Polyoxometalate by Enclosure in a Polyhedral Silver–Alkynyl Cage, *J. Am. Chem. Soc.* 131 (2009) 18257–18259.
- [27] Y.-P. Xie, T.C.W. Mak, Silver(I)-Ethyne Clusters Constructed with Phosphonate-Functionized Polyoxovanadates, *J. Am. Chem. Soc.* 133 (2011) 3760–3763.
- [28] S.C.K. Hau, P.-S. Cheng, T.C.W. Mak, Enlargement of Globular Silver Alkynide Cluster via Core Transformation, *J. Am. Chem. Soc.* 134 (2012) 2922–2925.
- [29] Y.-P. Xie, T.C.W. Mak, A pyrovanadate-templated silver(I) – ethynide cluster circumscribed by macrocyclic polyoxovanadate(V), *Chem. Commun.* (2012) 1123–1125.
- [30] Y.-P. Xie, T.C.W. Mak, High-Nuclearity Silver Ethynide Clusters Assembled with Phosphonate and Metavanadate Precursors, *Angew. Chem. Int. Ed.* 51 (2012) 8783–8786.
- [31] Y.-P. Xie, T.C.W. Mak, Silver-Ethyne Clusters with Oxovanadate Components, *J. Cluster Sci.* 23 (2012) 727–736.
- [32] K.A. Al-Farhan, M. H. Ja'far, O. M. Abu-Salah, The synthesis, identification, and X-ray structure of a novel cationic alkynyl silver cluster polymer $[\text{Ag}_3(\text{tBuC}\equiv\text{C})_2^+]_n$, *J. Organomet. Chem.* 579 (1999) 59–62.
- [33] HL (0.023 g, 0.010 mmol) was dissolved in CH_3OH (4 mL). Then $\text{tBuC}\equiv\text{CAg}$ (0.095 g, 0.500 mmol) and AgCF_3CO_2 (0.022 g, 0.100 mmol) were added under stirring. After 30 min, a clear solution was collected by filtration. Slow evaporation of the clear solution afforded colorless prismatic crystals **1** in *ca.* 16%. Elemental analysis (%) calcd for $\text{C}_{43}\text{H}_{49}\text{Ag}_7\text{F}_6\text{O}_8$: C 33.04, H 3.16; found: C 33.35, H 3.31; Selected IR data (KBr): 2005 ($\text{C}\equiv\text{C}$).
- [34] Crystallographic data. Complex **1**: monoclinic, $a = 13.625(1)$, $b = 29.180(3)$, $c = 13.767(1)$ Å, $\beta = 111.262(1)$ V = 5101(8) Å³, $T = 293$ K, space group $P2_1/c$, $Z = 4$, $\lambda = 0.71073$ Å, $\rho = 2.034$ cm⁻³, $\mu(\text{MoK}\alpha) = 2.699$ mm⁻¹, $R_1 = 0.0462$, $wR_2 = 0.1366$ for $I >$

$2\sigma(I)$, GOF = 1.117. Crystal data were collected on a Bruker Smart Apex II CCD diffractometer with Mo-K α radiation ($\lambda = 0.71073 \text{ \AA}$). The structures were solved by direct methods with SHELXS-97 and refined by full-matrix least-squares techniques using the SHELXL-97 program [36] within WINGX [37].

- [35] F. Migliaccio, A.W. Galston, On the Nature and Origin of the Calcium Asymmetry Arising during Gravitropic Response in Etiolated Pea Epicotyls, *Plant. Physiol.*, 85 (1987) 542-547.
- [36] G.M. Sheldrick, SHELXL-97, Program for Crystal Structure Solution; University of Göttingen: Göttingen, Germany, (1997).
- [37] L.J. Farrugia, WINGX, A Windows Program for Crystal Structure Analysis, University of Glasgow, UK, (1988).

Legends for figures


Fig. 1. Coordination environment of silver(I) atoms surrounded by four independent *tert*-butylethyne, one 9-hydroxy-9-fluorene-carboxylate, and two trifluoroacetate ligands in $\{[\text{Ag}_{14}(\text{C}\equiv\text{C}^t\text{Bu})_8(\text{L})_2(\text{CF}_3\text{CO}_2)_4]\cdot 2\text{CH}_3\text{OH}\}_n$ (**1**). The C \equiv C triple bond is represented by a bold rod. The Ag–C bonds are indicated by broken lines. All hydrogen atoms and methanol molecules have been omitted for clarity. Color code: Ag, violet; C, gray; O, red; F, lime. Ag1A–C31 = 2.325(8), Ag1–C15 = 2.334(9), Ag1–C16 = 2.647(9), Ag1A–C32 = 2.679(7), Ag2A–C37 = 2.349(9), Ag2–C23 = 2.521(9), Ag3–C23 = 2.134(7), Ag3A–C15 = 2.151(8), Ag4–C37 = 2.429(7), Ag4–C15 = 2.438(9), Ag4–C23 = 2.519(6), Ag5–C37 = 2.150(8), Ag5–C31 = 2.195(7), Ag6–C23 = 2.376(8), Ag6–C31 = 2.494(7), Ag6–C24 = 2.545(8), Ag7B–C31 = 2.256(8), Ag7B–C32 = 2.665(7). Symmetry code: A $1-x, -y, -z$; B $2-x, -y, -z$.

Fig. 2. (a) The centrosymmetric $[\text{Ag}_{14}(\text{C}\equiv\text{C}^t\text{Bu})_8(\text{CF}_3\text{CO}_2)_2]^{4+}$ cluster skeleton in complex **1**. H atoms are omitted for clarity. The carbon atoms of the ethynide group are represented as small black balls, and their bonds to silver atoms are indicated by broken lines. Atoms Ag7B and Ag7C are symmetry equivalents of Ag7. (b) Perspective view of the centre Ag₁₂ portion in $[\text{Ag}_{14}(\text{C}\equiv\text{C}^t\text{Bu})_8(\text{CF}_3\text{CO}_2)_2]^{4+}$ cluster skeleton. Color code: Ag, violet; C, gray; O, red; F, lime. Ag1 \cdots Ag5A = 2.899(2), Ag1 \cdots Ag3A = 2.985(2), Ag1 \cdots Ag2 = 3.076 (2), Ag2 \cdots Ag3 =

2.945(2), Ag2...Ag3A = 3.064(3), Ag2...Ag5A = 3.115(3), Ag2...Ag4 = 3.317(2), Ag3...Ag4 = 2.983(2), Ag3...Ag4A = 2.995(2), Ag3...Ag6 = 3.278(3), Ag4...Ag5 = 3.000(2), Ag4...Ag6 = 3.231(2), Ag5...Ag6 = 2.938(3), Ag5...Ag7B = 3.325(2). Symmetry code: A $1-x, -y, -z$; B $-1+x, y, z$; C $2-x, -y, -z$.

Fig. 3. Coordination polymeric chain in complex **1**. Each C≡C triple bond is represented by a bold rod. The Ag–C bonds are indicated by broken lines. All hydrogen atoms and methanol molecules have been omitted for clarity. Color code: Ag, violet; C, gray; O, red; F, lime.

Fig. 4. Packing of coordination polymeric chains in the crystal structure of **1** viewed parallel to the *a* axis. All methanol molecules have been omitted for clarity. Color code: Ag, violet; C, gray; O, red; F, lime.

Graphical abstract (pictogram)

ACCEPTED MANUSCRIPT

Graphical abstract (synopsis)

The crystalline silver(I)-ethynide complex $\{[Ag_{14}(C\equiv C^tBu)_8(L)_2(CF_3CO_2)_4]\cdot 2CH_3OH\}_n$ (HL = 9-hydroxy-9-fluorenicarboxylic acid) is composed of a packing of organosilver(I) coordination polymeric chains each containing centrosymmetric, bar-like $[Ag_{14}(C\equiv C^tBu)_8(CF_3CO_2)_2]^{4+}$ cluster segments bridged by pairs of L and trifluoroacetate ligands.

Highlights

- ▶ New silver(I)-ethynide complex featuring an infinite coordination polymeric chain.
- ▶ $[\text{Ag}_{14}(\text{C}\equiv\text{C}^t\text{Bu})_8(\text{CF}_3\text{CO}_2)_2]^{4+}$ cluster segments bridged by two different carboxylates.
- ▶ Versatile supramolecular synthon $\text{R}-\text{C}\equiv\text{C}\supset\text{Ag}_n$ for synthesis of organosilver complexes.




Fig. 1.




Fig. 2.




Fig. 3.

ACCEPTED MAN




Fig. 4.